

ms
Jazz The

The disease denoted non-A-non-B hepatitis is caused in many cases by the hepatitis C virus (HCV). HCV is a single-stranded encapsulated RNA virus whose genome is composed of about 9 to 10,000 bases. This genome codes for structural proteins (core and envelope proteins) as well as for non-structural proteins. The non-structural protein 3 (NS3) region of HCV contains a protease and a helicase. The protease activity is localized in the amino terminal third of the NS3 region.

EP-A-0 450 931¹ incorporated by reference discloses the complete nucleotide and amino acid sequence of HCV. In addition a combination of synthetic HCV antigens is described comprising a first HCV antigen from the C domain and at least one further HCV antigen from one of the non-structural domains NS3, NS4 or NS5 and the envelope domain S. A preferred antigen from the domain NS3 is an antigen denoted C33c which comprises ~~the~~ amino acids 1192 to 1457² encoded by ~~of~~ the HCV genome shown in Figure 1 of EP-A-0 450 931.

The international Patent Application W092/11370 concerns the cloning and sequencing of various polypeptides from the genome of HCV and the use of these polypeptides without any foreign protein parts in test kits and as a vaccine. A clone NS-3 deposited in connection with this application at the "Deutsche Sammlung für Mikroorganismen und Zellkulturen GmbH, Mascheroder Weg 1b, 38124 Braunschweig, BRD (DSMZ)" under the number 6847 contains the genetic information for a polypeptide of 527 amino acids from the NS3 region of HCV.

Mori et al., (Jpn. J. Cancer Res. 83 (1992), 264-268) describes the diagnostic detection of a HCV infection by determining viral antibodies in blood using viral proteins as antigens. These HCV proteins are expressed in E. coli as a fusion protein with β -galactosidase. A protein from the NS3 region containing the amino acids 1295 to 1541 of the HCV genome showed the highest sensitivity. However, a disadvantage of such fusion proteins is that cross-reactions with the fused protein part can occur which reduce the specificity of the test reaction.

A test for HCV in blood requires a high ^{degree of} specificity and sensitivity ~~of the test~~. Furthermore the antigen used for the test should be capable of expression in high yield and be stable. Previously known antigens from the HCV genome have disadvantages since they do not fulfil one or several of the above requirements.

Thus an object of the present invention ~~was~~ to provide a polypeptide, ^{encoded by} ~~from~~ the HCV genome in which these disadvantages of the state of the art are at least partially liminated and which, especially in comparison

22

a
a
a
a

NO: 1

SEQ ID NO. 1 shows the amino acid sequence of a polypeptide according to the invention. Amino acids 1 - 13 are ^{foreign}~~foreign~~ amino acids. Amino acids 14 - 295 originate from HCV. The polypeptide according to the invention particularly preferably contains ~~the~~ amino acids 14 - 295 of the amino acid sequence shown in SEQ ID NO. 1 and 2 or an amino acid sequence which is at

least 90 % homologous thereto.

09896032 062901
T06290 2E99860

The present invention also concerns a polypeptide defined as above which contains at least one marker group. All known marker groups come into consideration as the marker group that can be detected in a test system i.e. directly or indirectly detectable marker groups. In this connection a directly detectable marker group is understood as a group which produces a directly detectable signal e.g. a radioactive group, an enzyme-group, a luminescence group, a metal complex etc.. On the other hand the marker group can also be an indirectly detectable group e.g. a biotin or hapten group that is detectable by reaction with a suitable binding partner (streptavidin, avidin or anti-hapten antibody) which in turn carries a signal-generating group. The marker group can be coupled in a known manner to the antigen for example via a bifunctional spacer. Such processes for coupling marker groups to peptide antigens are known to a person skilled in the area of immunology and do not need to be described in detail here.

The polypeptide according to the invention preferably contains 1 to 12 and particularly preferably 3 to 7 marker groups.

a In addition, the invention concerns a polypeptide as defined above in which one or several of the sulfhydryl groups ^{of} ~~originating from~~ cysteine residues are present in a covalently modified form. Examples of suitable covalent modifying groups are maleimidodioxaoctylamine - (MADOO), N-methyl-maleinimide (NMM), iodoacetic acid and iodoacetamide. The covalent cysteine modification

a results in ~~a~~ particularly high specific immunological reactivity.

Directly or indirectly detectable marker groups are *a* ~~particularly~~ preferably covalently coupled to the sulfhydryl groups of the polypeptide. Examples of SH-reactive bifunctional linkers for coupling to sulfhydryl groups are maleimidopropylamine (MP), maleimidoethylamine (MEA) and maleimidodioxaoctylamine (MADOO).

a In addition, it may be preferable to use a polypeptide according to the invention in which one or several cysteine residues are replaced by other natural or artificial amino acids. Cysteine residues are preferably replaced by structurally analogous α -amino acids e.g. serine or α -aminobutyric acid. Cysteine substitutions *a* lead to ~~a~~ particularly high stability.

The polypeptide according to the invention has surprising advantages over already known polypeptides. Compared to the antigen C33 described in EP-A-0 450 931 which contains the amino acid residues 1192 to 1457 of the HCV sequence, the ^{polypeptides} ~~polypeptide~~ according to the invention ^{exhibits} ~~exhibits~~ a substantially higher specificity which is manifested in a statistically significant lower number of false positive results in negative sera. Compared to the antigen NS3 described in WO92/11370 which contains the region of amino acids 1007 to 1534 of the HCV sequence, the polypeptide according to the *a* invention has ~~a~~ considerably higher stability under test conditions. Compared to a polypeptide which contains amino acids 1227 to 1528 from the NS3 region of HCV, the polypeptide according to the invention has the advantage *a* of ~~an~~ improved expression efficiency and a higher

09896032-062901

a sensitivity. Due to these advantages the ~~polypeptide~~^{polypeptides}
a according to the invention ~~is~~^{are} substantially superior to
all previously known HCV antigens from the NS3 region.

a In addition, the invention concerns ~~a~~^{an isolated} nucleic acid which
codes for a polypeptide according to the invention. A
preferred example of such a nucleic acid is a foreign
a DNA, ~~insertion~~^{encoding the peptide of interest inserted} in the vector pUC-D26.

a SEQ ID ~~NO. 1~~^{NO:1} also shows the nucleotide sequence of a
nucleic acid according to the invention which codes for
a the ~~polypeptide~~^{polypeptides} of SEQ ID ~~NO. 1~~^{NO:1} and 2. Nucleotides 40 -
885 code for the region of the polypeptide derived from
HCV. The nucleic acid according to the invention
preferably contains (a) nucleotides 40 - 885 of the
nucleotide sequence shown in SEQ ID NO.1 or (b) a
nucleotide sequence that corresponds to a sequence from
(a) within the scope of the degeneracy of the genetic
code.

a The present invention, ~~in addition~~^{also} concerns a vector
which contains at least one copy of a nucleic acid
according to the invention. The vector according to the
invention is preferably a prokaryotic vector i.e. a
vector suitable for propagation in a prokaryotic host
cell. Examples of such vectors are shown in Sambrook et
al. (Molecular Cloning. A Laboratory Manual, 2nd Ed.,
Cold Spring Harbor Laboratory Press (1989)), especially
in chapters 1 to 4 and 17. The vector according to the
a invention is ~~particularly~~ preferably a circular plasmid.
The nucleic acid according to the invention is
preferably present on the vector under the control of a
promoter sequence which allows expression of the
polypeptide according to the invention. A preferred

09896032-062901

exempl of a vector according to the invention is pUC-D26.

Furthermore the invention concerns a cell which is transformed with at least one copy of a nucleic acid according to the invention or with a vector according to the invention. The cell is preferably a prokaryotic cell, ^{more} particularly preferably a gram-negative prokaryotic cell and most preferably an E. coli cell.

The polypeptide according to the invention is preferably used as an antigen in an immunological test procedure. On the other hand the polypeptide can, however, also be used as a helicase protein and, due to its excellent antigenic action, for the production of a vaccine against ~~a~~ HCV infection.

The present invention additionally concerns a method for the immunological determination of an antibody directed towards a hepatitis C virus in a sample liquid which is characterized in that the sample liquid is incubated with at least one polypeptide according to the invention and the antibody is detected via binding to the polypeptide. This immunological method of determination can be carried out using any known test format e.g. in a homogeneous immunoassay with a single reaction phase or in a heterogeneous immunoassay with more than one reaction phase. A heterogeneous test format is preferably used in which the presence of the antibody is detected in the presence of a reactive solid phase.

One embodiment of this test format is the so-called double antigen bridge test concept. In such a method the sample liquid is incubated with at least two

polypeptides P_1 and P_2 according to the invention in which polypeptide P_1 (a) is bound to a solid phase or (b) is present in a form capable of binding to a solid phase and polypeptide P_2 carries a marker group. The antibody in the sample liquid is detected by determining the label in the solid phase or/and in the liquid phase, preferably in the solid phase, by means of an immobilized complex i.e. one that is bound to the solid phase.

The test procedure preferably comprises mixing the sample liquid with a purified ^{labeled} antigen P_2 and with the purified solid phase antigen P_1 in order to obtain a ^{labeled} immobilized complex consisting of ^{labeled} antigen, antibody and solid phase-bound antigen. Compared to other test formats for detecting antibodies, the bridge test format leads to an improvement of the sensitivity ^{of the assay}, i.e. additional immunoglobulin classes such as IgM are recognized, and of the specificity, i.e., unspecific reactivities with anti-IgG conjugate are reduced.

The labelled antigen P_2 carries a directly or indirectly detectable marker group as described above. The solid phase antigen P_1 can ^{be bound, for example} ~~for example be bound~~ directly to the solid phase via a bifunctional spacer. However, P_1 is preferably a conjugate present in the solid phase of a polypeptide according to the invention and of a reaction partner of a specific binding system. The other reaction partner of the specific binding system is present bound to the solid phase. Examples of such specific binding systems are biotin/avidin, biotin/streptavidin, biotin/antibiotin, hapten/antihapten, carbohydrate/lectin and antibody or antibody fragment and antibody against this antibody or

TOPSECRET-062901

against the antibody fragment. The antigen P_1 is preferably in the form of a biotin conjugate.

In such a double antigen bridge test the polypeptide antigen according to the invention is preferably used in a soluble form in order to avoid increases in the blank_Λ value and an unfavourable signal/noise ratio due to aggregations of the antigen. For this purpose the antigen is already either expressed in a soluble form in a suitable expression system, or after expression in soluble form, it is renatured in vitro in a known manner. Furthermore in order to avoid the formation of covalently cross-linked molecular aggregates, the immunological test can be carried out under mild reducing conditions (addition of mild reducing reagents, preferably of sulfhydryl reagents, preferably DTT (dithiothreitol) or DTE (~~dithioerythritol~~ ^{dithioerythritol}) in a concentration range of 1 mmol/l to 25 mmol/l) or/and preferably an antigen with covalently modified sulfhydryl groups or/and an antigen with at least partially substituted cysteine residues is used.

The polypeptide antigen according to the invention is preferably used in a soluble form in such a double antigen bridge test in order to avoid increases in the blank and an unfavourable signal/noise ratio due to aggregations of the antigen. A detailed description of the bridge test format is given in EP-A-0 280 211.

~~the bridge test format is given in EPA 620-211.~~
a This patent application is incorporated by reference.
~~Reference is herewith made to this disclosure.~~

The polypeptide according to the invention can, however, also be used in other test formats. An example of this is an indirect immunoassay for recognizing a specific immunoglobulin by binding to an immobilized specific

antigen and indirect detection via a conjugate with a second antibody. In this embodiment of the method according to the invention the sample liquid is incubated with a polypeptide P_1 which is (a) bound to a solid phase or (b) is present in a form capable of binding to a solid phase and with a further antibody directed towards P_1 which carries a marker group. The antibody to be determined is detected indirectly by determination of the label in the solid phase ^{and/or} ~~or~~ and in the liquid phase, preferably in the solid phase. In this method the signal produced on the solid phase by an immobilized complex of labelled antibody and solid phase-bound antigen is indirectly proportional to the concentration of the antibodies to be determined in the sample liquid.

^{also}
a The present invention ~~in addition~~ concerns a reagent for the immunological determination of an antibody directed towards hepatitis C virus which contains at least one polypeptide according to the invention. If the reagent is used in a double antigen bridge test, it preferably contains at least two polypeptides P_1 and P_2 wherein polypeptide P_1 is (a) bound to a solid phase or (b) is present in a form capable of binding to a solid phase and polypeptide P_2 carries a marker group. Binding of the polypeptide P_1 to the solid phase can either be achieved by direct binding or by means of a specific binding pair, preferably streptavidin/avidin and biotin. The polypeptide P_1 is particularly preferably present in a biotinylated form.

If it is used in an indirect immunological test the reagent according to the invention preferably contains a polypeptide P_1 which is (a) bound to a solid phase or (b) is present in a form capable of binding to a solid

09896032-062901

A further area of application of the polypeptide according to the invention is to produce vaccines. For this the polypeptides according to the invention are preferably produced in a purified form and then brought into the form of injectable liquids which can either be solutions or suspensions of the polypeptides. The polypeptides can also be enclosed in liposomes. Further constituents of these vaccines ^{may include} ~~are for example~~ water, salt solutions, glucose or glycerol. In addition the vaccines ^{may} contain small amounts of auxiliary substances such as emulsifiers, buffer substances, and if necessary adjuvants which increase the immune response. The vaccines are usually administered parenterally by injection, preferably subcutaneously or intramuscularly.

Yet a further object of the present invention is a method for the immunological determination of an antibody directed towards a hepatitis C virus in a sample liquid in which the sample liquid is incubated with at least one polypeptide that contains sequence regions from the hepatitis C virus, in particular from the NS3 region of the hepatitis C virus, and the antibody is detected by binding to the polypeptide,

which is characterized in that one uses a polypeptide from a region which contains at least one cysteine residue and (a) the antibody is determined under reducing conditions, (b) one or several cysteine residues are covalently modified or/and (c) one or several cysteine residues are replaced by other amino acids.

In contrast to other antigenic regions from HCV the NS3 region contains a particularly large accumulation of cysteine residues . Although the NS3 protein synthesized during the course of virus replication is stable, the use of NS3 antigens under physiological buffer conditions, e.g. in immunological test procedures, has proven to be extremely problematic since the free sulfhydryl groups of the cysteine residues easily oxidize under these conditions. This leads to intramolecular as well as intermolecular cross-linking of the antigen which significantly reduces its immunological reactivity.

Surprisingly it was possible to significantly improve the immunological reactivity of these NS3 antigens by means of a test format in which one or several NS3 antigens with modified cysteine residues or/and with substituted cysteine residues are used or mild reducing conditions are present for example by addition of a sulfhydryl reagent. In this way an early recognition of seroconversion is achieved as well as a significant amplification of the measurement signal.

The present invention is described further by the following examples and sequence protocols.

09896032-062901

SEQ ID NO. 1: shows the nucleotide sequence of a preferred polypeptide according to the invention,
SEQ ID NO. 2: shows the amino acid sequence of a preferred nucleic acid according to the invention,
SEQ ID NO. 3: shows the nucleotide sequence of primer (1)
SEQ ID NO. 4: shows the nucleotide sequence of primer (2)
SEQ ID NO. 5: shows the nucleotide sequence of primer (3)
SEQ ID NO. 6: shows the nucleotide sequence of primer (4)
SEQ ID NO. 7: shows the nucleotide sequence of primer (5) and
SEQ ID NO. 8: shows the nucleotide sequence of primer (6).

EXAMPLES

Example 1

Cloning and expression of a polypeptide with amino acids 1207 to 1488 from the NS3 region of the hepatitis C virus

A DNA fragment was amplified by means of PCR starting with the clone NS3 (DSM 6847) using primers (1) and (2) whose nucleotide sequence is shown in SEQ ID NO. 3 and SEQ ID NO. 4. Sequences for cloning (BamHI, BspHI, EcoRI restriction cleavage sites) are located at the 5' end of this DNA fragment as well as an ATG codon and an AAA(Lys) codon to increase expression. Restriction sites for HindIII and EcoRI and a stop codon (^{TAA}~~TTA~~) are present at the 3' end. In primers (1) and (2) the region homologous to HCV starts at nucleotide No. 19 .

The DNA fragment obtained in this way was used in a pUC8 vector cleaved with BamHI and HindIII. The resulting plasmid was denoted pUC-D26.

106290-28095860

An E. coli strain JM109 (Yanisch-Perron et al., Gene 33 (1985), 103) transformed with plasmid pUC-D26 was incubated overnight in 100 ml medium (L broth/ampicillin). Next morning the culture was diluted in a 3 l flask with 900 ml 2x L-broth (10 g tryptone, 10 g yeast extract, 5 g NaCl per liter)/ampicillin. After addition of 2 ml glycerol and 1 to 2 drops silicon anti-foam emulsion (Serva Company) the culture was then shaken at ca. 185 rpm and 37°C for 2 hours. Induction and antigen production were achieved by adding 2 mmol/l of the inducer isopropylthio- β -D-galactoside (IPTG) and shaking further for 3 to 4 hours. Subsequently the bacteria were pelleted by centrifugation and processed further.

a The bacterial pellets from two ^{one liter} ~~1 l~~ cultures were resuspended in 200 ml 50 mmol/l Tris-HCl, pH 8.5, 0.2 mg/ml lysozyme and 2 mmol/l dithioerythritol (DTE). Subsequently EDTA (final concentration: 15 mmol/l), phenylmethylsulfonyl fluoride (final concentration: 1 mmol/l) and 4 mg DNase were added. The suspension was mixed for several minutes with a magnetic stirrer and incubated for 45 minutes at 37°C in a water bath.

Afterwards Triton-X100 (final concentration: 1 %) was added and it was stirred for 30 minutes in a magnetic stirrer. After freezing at -20°C overnight and thawing, the cells were stirred for at least 1 hour at 37°C and sonified if necessary. The incubation at 37°C and/or ultrasonic treatment should be continued until the viscosity of the suspension of lysed cells has decreased significantly.

09896032 062901

Subsequently, ^{this} it was centrifuged for 20 minutes at 35,000 g and 4°C. The resulting pellet was resuspended in 30 ml 50 mmol/l Tris-HCl, pH 8.5, 2 mmol/l DTE, 150 mmol/l EDTA and 1.5 % OGP (octyl-β-D-glucopyranoside, Biomol Company). This suspension was stirred vigorously at room temperature with a magnetic stirrer for at least three hours and subsequently centrifuged at 35,000 g and 4°C for 20 minutes.

- a The pellet was dissolved in 100 ml ^{of} 8 mol/l urea, 20 mmol/l Tris-HCl, pH 8.5, 2 mmol/l DTE and stirred.
- a The antigen which was now dissolved, can be frozen at -20°C until further processing.

The protein was purified by the chromatographic steps described ⁱⁿ ~~in the following~~ which were carried out at room temperature. The antigen was stored between each of the chromatographic steps at -20°C.

The first chromatographic step was carried out on a Q-Sepharose Fast Flow column (Pharmacia) with a 20 mmol/l Tris-HCl pH 8.5, 8 mol/l urea, 2 mmol/l DTE buffer. It was eluted with a NaCl gradient (0 to 0.7 mol/l). The void volume and the fractions were tested by means of SDS PAGE. The polypeptide according to the invention is located in the first main peak. The positive fractions were pooled and dialyzed overnight against a 10-fold volume of 4 mol/l urea, 2 mmol/l DTE, 20 mmol/l Tris-HCl pH 7.3.

The same column was used in the second chromatographic step. The column buffer was the dialysis buffer used after the first chromatographic step. It was eluted using a NaCl gradient (0 to 0.5 mol/l) and subsequently

09896033-062901

with 1 mol/l NaCl, NaOH pH 13, 4 mol/l urea.

The positive fractions were pooled and dialyzed overnight against the same buffer as above.

Finally it was chromatographed on a S-Sepharose Fast Flow column (Pharmacia). The buffer and elution conditions were the same as in the previous chromatographic step.

a The antigen has a size of ca. 41 kDa, ^{on}~~in the~~ SDS polyacrylamide gel. The yield was ca. 10 mg antigen per litre culture medium.

EXAMPLE 2

Expression of the antigen according to the invention in comparison with other antigens from the HCV NS3 region

The following antigens were expressed:

- a) antigen D26 according to the invention (amino acids 1207 to 1488)
- b) antigen C33 (amino acids 1192 to 1457 according to EP-A 0 450 931)
- c) antigen D27 (amino acids 1227 to 1528)
- d) antigen NS3 (amino acids 1007-1534)

The antigen NS3 was expressed using the deposited clone NS-3 (DSM 6847). Cloning and expression of the antigens C33 and D27 were carried out according to the method described in example 1. The coding regions of amino acids 1192 to 1457 for C33 and amino acids 1227 to 1528 for D27 were amplified by standard methods using PCR starting with the plasmid pUC-N3 from the clone NS-3.

09896032-DE2901

The primers (3) and (4) were used for C33 whose nucleotide sequences are shown in SEQ ID NO.5 and SEQ ID NO. 6. The primers (5) and (6) were used for D27 whose nucleotide sequences are shown in SEQ ID NO.7 and SEQ ID NO. 8. The region homologous to HCV starts at nucleotide No. 19 in primers (3) and (5) and at nucleotide No. 13 in primers (4) and (6).

After treatment with the restriction enzymes BamHI and HindIII and subsequent purification by agarose gel electrophoresis the amplified DNA fragments were inserted into the vector pUC8 that had been cleaved with BamHI and HindIII. Both antigens expressed by the pUC vectors have a region of 13 non HCV-coded foreign amino acids at the N-terminal end (Met-Thr-Met-Ile-Thr-Asn-Ser-Arg-Gly-Ser-Ile-Met-Lys).

Afterwards the plasmids were transformed in E. coli JM109. Clones which receive a DNA fragment coding for the C33 antigen express an antigen with a molecular weight of ca. 34 kDa (SDS-PAGE). In clones with a DNA fragment that codes for antigen D27, a band of 48 kDa is found.

In the case of C33 its percentage of the total protein is ca. 10 % and is about the same magnitude as the polypeptide D26 according to the invention. A considerably smaller expression in the range of 5 % or less is found for D27.

T06290"22096860

Lysates of clones expressing D26, D27 and C33 were concurrently separated in an SDS gel and transferred onto nitrocellulose. After incubating overnight with various HCV-positive sera in a 1:100 dilution, bound antibodies were detected by means of an anti-human IgG antibody-peroxidase conjugate and a colour reaction with 3,3'-diaminobenzidine tetrachloride (DAB)/hydrogen peroxide.

^{Good}
~~a good~~ reactivity is found in all HCV antigen segments with strongly positive sera. In the case of weakly positive NS3-HCV sera the C33 antigen exhibits in some cases the same and in other cases a slightly weaker reactivity in comparison with the antigen according to the invention. However, in the case of D27 a considerably reduced colour reaction with weak NS3-positive sera is found.

EXAMPLE 3

Investigation of the specificity and sensitivity of various antigens from the NS3 region of HCV

The reactivity of various antigens from the NS3 region was comparatively assessed on a total of 960 negative sera using the indirect test concept. In this evaluation the antigen D26 according to the invention was found to have a significantly superior specificity compared to the reference antigen C33 which manifests itself as a significantly reduced number of false positive results.

09696032 "062901

Construct	false positive classification	correct negative classification
C33	8	952
D26	2	958

In addition the reactivity of various NS3 constructs was comparatively assessed using the indirect test concept on 20 sera of proven HCV status. A reduced sensitivity of antigen D27 was found with comparable sensitivity of the constructs NS3 (amino acids 1007 to 1534), C33, D26 and D27.

Construct	correct positive classification	false negative classification
NS3	20	0
C33	20	0
D26	20	0
D27	19	1

EXAMPLE 4

Investigation of the stability of antigens from the NS3 region of HCV

The stability of the antigens NS3 and D26 was comparatively evaluated using the indirect test concept. Antigen NS3 was found to have a significantly poorer stability compared to antigen D26 according to the invention. The stability of the antigens was investigated after 72 hours incubation at 37°C.

Construct	Sample No.	Signal recovery
NS3	1	< 20 %
	2	< 20 %
D26	1	99 %
	2	103 %

EXAMPLE 5

Investigation of the reactivity of the HCV-NS3 helicase antigen in the presence or absence of reducing reagents

The time of seroconversion was tested on the bases of the reactivity of serum samples collected at various times with the HCV-NS3 helicase antigen under physiological conditions with or without 20 mmol/l DTT. The test concept was a double-antigen bridge test for the class-independent recognition of all immunoglobulins in which an antigen provided with an electrochemical marker group (ruthenium metal complex) and an antigen bindable to a solid phase (biotinylated antigen) was used.

The results of this test are shown in the following table. It can be seen that it was possible to recognize seroconversion 38 days earlier in the presence of 20 mmol/l DTT. In addition an improved signal strength was found in the presence of DTT.

106290" 2E096860

TABLE

Day of blood collection	Days after start of blood collection	Reactivity of the NS3 antigen without DTT (signal/cut off)	Reactivity of the NS3 antigen with DTT (signal/cut off)
28.07.1988	0	0.1	0.1
01.08.1988	4	0.1	0.1
08.08.1988	11	0.1	0.1
11.08.1988	14	0.1	0.1
15.08.1988	18	0.1	0.1
25.08.1988	28	0.1	0.1
29.08.1988	32	0.1	1.6*
14.09.1988	48	0.1	6.5*
05.10.1988	69	1.3*	4.8*
19.10.1988	83	2.1*	6.8*

* positive signal

09896032-052901